CHEMDRAW: INSIDER SECRETS EVERY CHEMIST SHOULD KNOW

JINSOO KIM, PHD, MANAGER, AMERICAS FAS, REVVITY SIGNALS

STUART CHALK, PHD, PROFESSOR, DEPARTMENT OF CHEMISTRY, UNIVERSITY OF NORTH FLORIDA

LEARN FROM THE BEST AND BRIGHTEST MINDS IN CHEMISTRY

How does ChemDraw transform end user workflow to maximize speed and efficiency?

This is primarily achieved by shifting from traditional menu-driven drawing to a keyboard-centric approach, mirroring the efficiency of shortcuts in applications like Microsoft Excel. This allows for the rapid creation of complex drawings. Dr. Kim demonstrated drawing Zorelto (a top-20 pharmaceutical) by effectively typing it out and constructing a multi-step reaction scheme in under a minute.

Key Hotkeys and Shortcuts

Feature	Keystroke(s)	Description & Use Case
Molecule Selection	Double-click on any atom or bond	Instantly selects the entire connected molecule. This is highly effective for deconvoluting and moving structures that are tightly packed or overlapping on the canvas.
Proportional Scaling	`Ctrl` + Click & Drag handle	Scales a structure while maintaining the correct proportions for bond thickness and heteroatom font size. This ensures high-fidelity graphics when enlarging for posters or shrinking for publications.
Precise Rotation	`Alt` + Up/Down Arrow `Alt` + Left/Right Arrow	Rotates the selected structure in 15-degree increments (Up/Down) or 1-degree increments (Left/Right), offering finer control than the mouse handle.
Structure Joining	`Ctrl` + `J`	Joins two selected atoms. The first atom selected remains stationary while the second moves to it. This feature circumvents automatic "structure cleanup," preserving specific user-drawn conformations (e.g., a "tent-shaped" indole) and can join atoms even if it violates valency (a warning is provided).

How has ChemDraw introduced a range of features that broadens the utility beyond standard organic chemistry drawing?

Alternative Structure Input Methods

To accommodate users with different workflows and expertise, ChemDraw supports multiple methods for adding chemical content to the canvas without manual drawing:

- **Name-to-Structure:** Generates a structure from its chemical name.
- **CAS Number:** Generates a structure from its CAS registry number.
- **Cheminformatics Strings:** Pasting a SMILES or InChI key string directly onto the canvas (Ctrl + V) automatically converts it into its 2D structure.
- **Biopolymer Sequences:** Using the "Paste Special" function, FASTA sequences for peptides, DNA, or RNA can be pasted to generate corresponding biopolymer representations.

Polymer Chemistry Enhancements

Addressing a historically "lightly served area," ChemDraw now allows polymer repeat units to be defined by Average Molecular Weight. By right-clicking a polymer bracket, a user can input a target weight (e.g., 2000 for PEG 2K), and the software calculates the average number of subunits required to achieve that weight.

On-Canvas 3D Visualization

- **Feature:** A "3D Cleanup" function (Ctrl + Shift + D) is now available directly on the 2D canvas.
- **Functionality:** It applies a force field to a selected structure, generating a more realistic, folded 3D representation without leaving the ChemDraw environment.
- **Purpose:** This provides a quick, convenient method for visualizing the 3D nature of a molecule. It is not intended as a full replacement for the more powerful computational tools in the separate Chem3D application but serves as an effective visualization aid.

CHEMDRAW: INSIDER SECRETS EVERY CHEMIST SHOULD KNOW

JINSOO KIM, PHD, MANAGER, AMERICAS FAS, REVVITY SIGNALS

STUART CHALK, PHD, PROFESSOR, DEPARTMENT OF CHEMISTRY, UNIVERSITY OF NORTH FLORIDA

LEARN FROM THE BEST AND BRIGHTEST MINDS IN CHEMISTRY

What tools does ChemDraw offer for users in patent analysis or library synthesis which will allow them to define and automatically generate large libraries from a single generic structure?

Defining Variable Structures

Users can create a comprehensive generic structure using a combination of notations:

- **Variable Heteroatoms:** Square brackets with comma-separated elements (e.g., [N,O,S]) define points where different heteroatoms can exist.
- **Expanding Rings:** Curly brackets (e.g., {CH2}1-3) define a variable number of repeating units, allowing for ring size variation.
- **Wandering Substituents:** A variable attachment point (represented by a star) can be linked to multiple atoms on a ring, denoting a substituent that can appear at different positions.
- **R-Group Tables:** An R-group on the core structure can be linked to a table defining a list of possible substituents.

Library Generation and Management

- **Enumeration:** The "Expand Generic Structure" command automatically draws all possible permutations of the defined variables. The presenter demonstrated generating an 81-member library in approximately five seconds.
- **Output Limit:** The software will display up to 200 enumerated structures directly on the canvas.
- **Large Libraries:** For libraries exceeding 200 compounds, the output can be sent directly to an SD file, a text-based format that can be opened in other software to access the full library (e.g., thousands of compounds).
- **Visual Analysis with Coloring:** By coloring specific fragments or R-groups in the initial generic structure, the colors are propagated through the enumeration. This allows for trivial visual identification of subsets within the generated library (e.g., all para-substituted analogs are highlighted in yellow).

What sophisticated toolset does the web-based ChemDraw+ offer specifically for biopolymer chemistry?

Biopolymer Representation and Libraries

- **Monomer Libraries:** Users can draw biopolymers using several built-in libraries:
  - Natural Monomers (amino acids, nucleotides)
  - Pistoia Alliance Standard Monomers
  - Revity Signals Curated Libraries
- **Custom Monomers:** Organizations can create and import their own libraries of custom monomers (e.g., Peptide Nucleic Acids - PNAs).
- **Mix-and-Match Capability:** It is possible to combine monomers from different libraries within the same sequence (e.g., attaching a PNA to a natural peptide).

Embedded Chemical Intelligence

The software performs complex chemical interpretations behind a simple user interface:

- **Hover-over Information:** Hovering over a single-letter monomer reveals its full chemical name and structure.
- **Complementary Strands:** The software can automatically generate the correct complementary strand for a DNA or RNA sequence, correctly substituting uracil for thymine where appropriate.
- **Hydrogen Bonding:** When drawing complementary strands, it correctly represents the two hydrogen bonds between A-U/T pairs and the three hydrogen bonds between C-G pairs.
- **Expandable Structures:** Sequences can be expanded to their full chemical structures to allow for side-chain modifications.

What are some future directions and ideas about AI integration for ChemDraw?

Revity is actively developing AI capabilities for ChemDraw with a phased approach:

- **Current/Near-Term:** The initial focus is on suggestive AI, where the software will propose reaction conditions (catalysts, solvents) for a user-defined transformation (reactants to products). This is currently in early testing.
- **Long-Term Vision:** The ultimate goal is to develop generative AI that can create entire reaction schemes from a high-level command, such as "Draw a Suzuki coupling followed by an esterification." This would represent a significant leap in automating chemical content creation.

LabNotes are AI-generated using speaker slides and audio from live ACS Webinars.

Scan or click for ACS Webinar Website!

ACS Chemistry for Life®

CHEMDRAW: INSIDER SECRETS EVERY CHEMIST SHOULD KNOW

JINSOO KIM, PHD, MANAGER, AMERICAS FAS, REVVITY SIGNALS

STUART CHALK, PHD, PROFESSOR, DEPARTMENT OF CHEMISTRY, UNIVERSITY OF NORTH FLORIDA

LEARN FROM THE BEST AND BRIGHTEST MINDS IN CHEMISTRY

GLOSSARY OF KEY TERMS

- **3D Cleanup:** A feature in ChemDraw, accessible via the icon or Ctrl+Shift+D, that applies a force field to a 2D structure to generate a quick, more realistic 3D representation directly on the canvas.
- **Annotation:** The process of adding notes or enhanced stereochemical information to structures.
- **Biopolymers:** Large biological molecules like peptides (amino acids) and oligonucleotides (DNA, RNA) that can be represented and manipulated in ChemDraw.
- **CAS Number:** A unique numerical identifier assigned to every chemical substance by the Chemical Abstracts Service. ChemDraw can generate a structure from an input CAS number.
- **CDX File:** The standard file format for documents created and saved in ChemDraw.
- **Chem3D:** A companion software to ChemDraw that provides more powerful and advanced 3D molecular modeling, visualization, and calculations.
- **ChemDraw:** A software program celebrating its 40th year, regarded as the de facto standard for digital chemical structure drawing.
- **ChemDraw for Excel:** An add-on for ChemDraw that allows users to create SD files and manage chemical structures and data within a spreadsheet environment.
- **ChemDraw Plus:** A browser-based, web-enabled version of ChemDraw that is part of the Signals ChemDraw package, allowing access and editing from any computer with a web browser.
- **Chem-informatics:** A scientific discipline that involves the use of computer and informational techniques applied to a range of problems in the field of chemistry.
- **Enumeration:** The process by which ChemDraw automatically draws out all possible individual structures from a defined generic structure.
- **FASTA:** A text-based format for representing nucleotide or peptide sequences. ChemDraw can interpret pasted FASTA text to generate corresponding biopolymer structures.
- **Generic Structures:** A feature that allows users to represent a large library of related compounds using a single core structure with defined variables, such as R-groups, expanding rings, and variable attachments.
- **HELM (Hierarchical Editing Language for Macromolecules):** A notation standard for representing complex biomolecules, supported by ChemDraw's biopolymer tools.
- **Hotkeys:** Single-key keyboard shortcuts (e.g., typing 'N' over an atom to create an amine) that allow users to draw and modify structures without using menus, greatly increasing efficiency.
- **InChI (Key):** The IUPAC International Chemical Identifier, a textual identifier for chemical substances. ChemDraw can generate structures from pasted InChI strings.
- **Join Feature:** A function (Ctrl+J) that creates a bond between two pre-selected atoms from different structures, moving the second structure to join the first while preserving the geometry of both fragments.
- **Pistoia Alliance:** An organization that has defined a stock library of common, non-natural monomers, which are available for use in ChemDraw's biopolymer drawing tools.
- **Polymer Brackets:** A drawing tool used to enclose a repeating monomer unit in a polymer. The properties can be defined by the number of repeats or by the average molecular weight of the polymer.
- **R-Group Table:** A table used in conjunction with generic structures where a user can define a list of different substituents for a specific R-group label (e.g., R1, R2) on the core structure.
- **Revity Signals:** The company that develops and manages ChemDraw and its associated products, such as Signals Notebook.
- **SD File (SDF):** Structure-Data File. A text-based file format that stores information about one or more molecules, allowing chemical structures and associated data to be exported and shared.
- **Signals ChemDraw:** The newest version of ChemDraw, which integrates the desktop application with a cloud-based system and the web-based ChemDraw Plus.
- **SMILES:** Simplified Molecular-Input Line-Entry System. A line notation for describing the structure of chemical species using short ASCII strings. ChemDraw can generate structures from pasted SMILES strings.
- **Stylesheet:** A set of predefined display settings (e.g., background color, bond color, font) that determines the appearance of the ChemDraw canvas. Users can select from presets like ACS 1996 or create their own.
- **Variable Attachment:** A feature in generic structures, represented by a star, that allows a substituent to be attached to multiple possible positions on a ring or structure.

LabNotes are AI-generated using speaker slides and audio from live ACS Webinars.

Scan or click for ACS Webinar Website!

ACS Chemistry for Life®